

Therapist Recertification Workshop

Workshop Title: Designing Personalized Freezing of Gait Interventions to Benefit Both FOG and Functional Mobility

Date

April 7, 2025

Location

Delivered via Zoom
Eastern Time

Workshop Fee

\$350 per person

Discounts available for groups of 2 or more



In this workshop you will learn to:

- Appreciate the heterogeneity and the complexity of FOG and how that impacts our ability to identify “freezers / non-freezers” and capture their response to our clinical interventions.
- Develop personalized proactive and rescue-focused interventions for clients with FOG that are informed by a theoretical understanding of the mechanisms that underlie FOG.

Who is eligible?

- Physical Therapists, Physical Therapy Assistants
- Occupational Therapists, Occupational Therapy Assistants
- PT, PTA, DPT, OT, OTA students

For additional information see the PWR! Workshop registration page



501(C)3 Non-Profit Organization

Are continuing education hours offered?

For details see the continuing education information page at <https://www.pwr4life.org/ceu-information/>

Earn 5-6 contact hours

My Time	Eastern Time	Topic
	11:00 am	Welcome and Housekeeping
	11:15 am	Freezing of Gait (FOG) Introduction <ul style="list-style-type: none"> • Terminology and Phenomenology • FOG / Festination vs. Non-Gait Freezing / Festination
	11:55 am	Mechanisms Underlying FOG and Implications for Intervention <ul style="list-style-type: none"> • Interference - Decreased reserve leads to crosstalk • Threshold - Gait instability • Decoupling - Posture with gait • Cognitive - Frontal executive dysfunction • Sensori-perceptual processing deficits
	1:20 pm	Break
	1:30 pm	Implications of FOG on Functional Mobility <ul style="list-style-type: none"> • Techniques and training tools for learning and retention • New paradigms of ongoing access to rehab and exercise for life
	1:50 pm	Video Case Study #1 <ul style="list-style-type: none"> • Assessment considerations and Level 3 goals for a <u>single</u> plan of care • Level 1-2-3 sample activities for addressing Level 3 goals
	3:00 pm	Break
	3:10 pm	Video Case Study #2 <ul style="list-style-type: none"> • Assessment considerations and changing Level 3 goals <u>over time</u> • Level 1-2-3 sample activities for addressing Level 3 goals <u>over time</u>
	4:15 pm	Q&A
	4:30 pm	End of PWR!Moves Therapist Freezing of Gait Recertification Workshop

Most people (~80%), if not all, persons with PD (PwPD) will experience freezing of gait (FOG) as the disease progresses into advanced stages. However, 21% to 27% of PwPD report FOG even in the early stages. This presentation will define the terminology and complex phenomenology underlying FOG. Participants will appreciate how this greatly interferes with research studies trying to clearly identify freezers from non-freezers so that we can better understand the data, the underlying mechanisms and the response to therapeutic interventions.

Participants will be introduced to screening tools and assessments that can be used for early identification or for therapy outcomes after FOG-focused interventions. We will review 5 theoretical hypotheses (models) to help understand the pathophysiology underlying FOG. These will include the categories of: Interference, Threshold, Decoupling, Cognitive or Sensori-perceptual. For each model, we will first discuss examples of behaviors that we may be able to observe or elicit in the clinic that support different models and how that can provide insight for designing interventions.

As we go through the models, we will introduce current research supporting interventions that may address one or more of these models. Emphasis will be on a recent meta-analysis that suggests that rehabilitation that targets multiple behavioral training modes related to 1) reducing the frequency or emergence of FOG episodes (proactive goals) and 2) circumventing and alleviating imminent FOG episodes (rescue goals) are efficacious. Altogether, these studies support the importance of starting early, incorporating cognitive strategies, challenging the intensity and complexity of training, targeting aerobic exercise to enhance neuroplasticity and addressing the unique challenges of sustained practice.

The severity of functional mobility impairment in PwP is affected by dopamine-dependent disturbances that alter cortico-striatal networks and interfere with the ability to express habitual-automatic actions in complex real-world scenarios. Therefore, each of the proposed mechanisms that underlie FOG; may also contribute to the loss of automaticity and the deterioration of functional mobility. In addition, freezing and festination can occur in non-gait functions that require repetition and/or coordination across limbs or multiple structures like speech, repetitive finger or hand movements, bilateral reach and grasp movements, and during upper or lower limb floor crawling movements.

For that reason, we propose that FOG interventions should consider the impact FOG has on ALL aspects of functional mobility. We will use Video case studies (n=2) to illustrate how FOG-specific interventions can be integrated within the PWR!Moves® Retrain Functional Mobility Framework allowing therapists to address personalized goals related to FOG but also everyday mobility and transfers in multiple positions or environments (home, work, recreation, community group exercise). Assessments used in the case studies, role of instruction, sensory feedback, use of simple multipurpose equipment and dosage consideration will be highlighted as part of case study discussion.

Upon successful completion of this workshop, participants will be recertified as PWR!Moves Certified Therapists for another three years!

Methods of Instruction

- Brief lectures with time for Q&A and response to chat.
- Pre-recorded video cases (n=2) showing therapists implementing the curriculum with people with PD of varying disease severity with freezing of gait.
- Faculty led group discussion to answer questions and review highlights for each video case study.

Course Objectives

- Develop personalized functional mobility interventions for clients with FOG that include FOG-specific motor and cognitive challenges.
- Interpret the proposed mechanisms that underlie FOG to inform your clinical reasoning for designing and prioritizing FOG proactive and rescue-focused goals and activities.
- Utilize assessments and interview techniques that may help characterize and capture FOG and non-gait freezing episodes for early identification, setting goals and demonstrating therapy outcomes.
- Use training tools related to cueing, instruction, feedback, aerobic priming, and simple equipment to optimize quality of movement and learning.
- Recognize the importance of dosage and sustained practice



Becky G. Farley, PT, MSPT, PhD

Dr. Becky Farley is a physical therapist, neuroscientist, Parkinson exercise specialist, Chief Scientific Officer and Founder of Parkinson Wellness Recovery | PWR!. She received a PhD in Neuroscience from the University of Arizona, a Master of Science in Physical Therapy from the University of North Carolina, and a Bachelor of Physical Therapy from the University of Oklahoma. She is a published author on exercise for people with Parkinson disease and gives public and medical seminars worldwide. Her postdoctoral research investigated the muscle activation deficits underlying bradykinesia in people with PD. She was awarded, and completed, an R21 NIH-funded randomized clinical trial to establish the benefits of LSVT BIG[®], the first whole-body, amplitude-focused, physical and occupational therapy exercise approach for individuals with PD. Dr. Farley also created PWR! Moves, a more flexible Parkinson-specific exercise approach that directly targets the training of amplitude into building blocks of function. Each building block counteracts a primary motor control deficit shown by research to interfere with everyday mobility. Dr. Farley has been training therapists and fitness professionals for the last 20 years and is now focusing on publishing data from the Tucson-based PWR!Gym[®] and integrating new research into PWR!Moves[®] workshops and PWR!Gym programs. She believes lifelong access to integrated rehabilitation and community exercise and wellness programming is necessary to optimize and perpetuate functional mobility benefits and to slow disease progression.



Jennifer Bazan-Wigle, PT, DPT, CEEAA[®]

Jennifer Bazan-Wigle has worked in neurological rehabilitation for the entirety of her physical therapy career. She is currently a physical therapist at Parkinson Wellness Recovery's PWR!Gym in Tucson, AZ, where she specializes in one-on-one rehabilitation and group exercise instruction with people with Parkinson disease. Since 2013, she has focused on honing her expertise in treating the movement disorder and Parkinson's population, emphasizing freezing of gait and advanced PD. Jennifer is a PWR!Moves Certified Therapist, PWR!Moves Certified Instructor, and a Certified Exercise Expert for the Aging Adult (CEEAA). Jennifer has delivered community, academic, and peer-reviewed presentations on Parkinson disease in the US and internationally. As an integral part of the PWR! Faculty, Jennifer has worked closely with Dr. Becky Farley to develop course content for PWR!Moves Therapist and Instructor Training and Certification Workshops, and has delivered over 100 continuing education workshops, across the US and world. In doing so, Jennifer has helped thousands of physical therapists, occupational therapists, and fitness professionals implement evidence-based rehabilitation and group exercise for people with Parkinson disease.



Shelley Hockensmith, PT, MPT

Shelley Hockensmith is a physical therapist with over 20 years of experience in outpatient neurological rehab settings. She graduated from the University of Evansville with her MPT in 2003 and in 2008 became a Board Certified Neurologic Clinical Specialist re-certifying in 2018. She has experience in private practice and hospital-based multi-disciplinary neurologic teams working with people with neurological disorders such as stroke, multiple sclerosis, brain injuries, spinal cord injuries, and movement disorders. She also was fortunate to work in a specialized vestibular and balance disorder clinic as both clinician and coordinator with a team of audiologists and physical therapists. As an avid believer in the power of exercise for people with Parkinson Disease, she became certified in LSVT BIG in 2007, attended one of the first PWR!Moves workshops, and eventually began working at the PWR!Gym in 2019 as a PWR! Moves[®] Certified Therapist. She joined the PWR!Moves[®] faculty in 2022.



Emily Borchers PT, DPT

Emily Borchers is a physical therapist with almost 10 years of experience working with people with Parkinson disease throughout her career. After graduating from the Ohio State University with her DPT in 2014, she began working as a physical therapist at the PWR!Gym® in Tucson, AZ where she developed a passion for helping people with Parkinson disease. In her 7 years of working at the PWR!Gym, she specialized in providing one-on-one rehabilitation and group exercise instruction for people with Parkinson disease, assisted with research conducted at the PWR!Gym including a peer-reviewed publication and

was Physical Therapy Manager where she learned how to navigate Medicare reimbursement issues to meet the ongoing rehabilitation needs of people with Parkinson disease. Emily also has experience working with people with Parkinson disease and other neurological conditions including stroke, brain injury, and spinal cord injury in the inpatient rehab setting as part of an interdisciplinary team. She now works at Banner Alzheimer's Institute in Tucson, AZ where she continues to develop her skills in working with people with Parkinson disease and other cognitive diseases including Alzheimer's and Lewy Body Dementia for outpatient rehabilitation services. She joined the PWR! Faculty in 2023. Emily is passionate about empowering people with Parkinson disease and implementing a proactive approach to ongoing rehabilitative and exercise services for improved quality of life.

1. Agosta F, Gatti R, Sarasso E, Volonté MA, Canu E, Meani A, Sarro L, Copetti M, Cattysse E, Kerckhofs E, Comi G, Falini A, Filippi M. Brain plasticity in Parkinson's disease with freezing of gait induced by action observation training. *J Neurol*. 2017 Jan;264(1):88-101. doi: 10.1007/s00415-016-8309-7. Epub 2016 Oct 24. PMID: 27778161.
2. Ahlskog JE. Common Myths and Misconceptions That Sidetrack Parkinson Disease Treatment, to the Detriment of Patients. *Mayo Clin Proc*. 2020;95(10):2225-2234. doi:10.1016/j.mayocp.2020.02.006
3. Ahlskog JE. Cheaper, simpler, and better: Tips for treating seniors with Parkinson disease. *Mayo Clin Proc*. 2011;86(12):1211-1216. doi:10.4065/mcp.2011.0443
4. Anke H, Snijders, Charlotte A, Haaxma, Yolien J, Hagen, Marten, Munneke, Bastiaan R, Bloem, Freezer or non-freezer: Clinical assessment of freezing of gait, *Parkinsonism & Related Disorders*, Volume 18, Issue 2, 2012, Pages 149-154
5. Beck EN, Ehgoetz Martens KA, Almeida QJ. Freezing of Gait in Parkinson's Disease: An Overload Problem? *PLoS One*. 2015 Dec 17;10(12):e0144986. doi: 10.1371/journal.pone.0144986. PMID: 26678262; PMCID: PMC4682987
6. Bekkers EMJ, Dijkstra BW, Dockx K, Heremans E, Verschueren SMP, Nieuwboer A. Clinical balance scales indicate worse postural control in people with Parkinson's disease who exhibit freezing of gait compared to those who do not: A meta-analysis. *Gait Posture*. 2017 Jul;56:134-140. doi: 10.1016/j.gaitpost.2017.05.009. Epub 2017 May 11. PMID: 28544951.
7. Bekkers EMJ, Dijkstra BW, Heremans E, Verschueren SMP, Bloem BR, Nieuwboer A. Balancing between the two: Are freezing of gait and postural instability in Parkinson's disease connected? *Neurosci Biobehav Rev*. 2018;94:113-125. doi:10.1016/j.neubiorev.2018.08.008
8. Bekkers EMJ, Mirelman A, Alcock L, Rochester L, Nieuwhof F, Bloem BR, Pelosin E, Avanzino L, Cereatti A, Della Croce U, Hausdorff JM, Nieuwboer A. Do Patients With Parkinson's Disease With Freezing of Gait Respond Differently Than Those Without to Treadmill Training Augmented by Virtual Reality? *Neurorehabil Neural Repair*. 2020 May;34(5):440-449
9. Bryant MS, Workman CD, Hou JG, Henson HK, York MK. Acute and Long-Term Effects of Multidirectional Treadmill Training on Gait and Balance in Parkinson Disease. *PM&R*. 2016 Dec;8(12):1151-1158. doi: 10.1016/j.pmrj.2016.05.001. Epub 2016 May 10. PMID: 27178378.
10. Chow R, Tripp BP, Rzonczinski D, Almeida QJ. Investigating Therapies for Freezing of Gait Targeting the Cognitive, Limbic, and Sensorimotor Domains. *Neurorehabil Neural Repair*. 2021;35(3):290-299. doi:10.1177/1545968321992331
11. Cosentino C, Baccini M, Putzolu M, Ristori D, Avanzino L, Pelosin E. Effectiveness of Physiotherapy on Freezing of Gait in Parkinson's Disease: A Systematic Review and Meta-Analyses. *Mov Disord*. 2020 Apr;35(4):523-536. doi: 10.1002/mds.27936. Epub 2019 Dec 4.
12. Delgado-Alvarado M, Marano M, Santurtún A, Urtiaga-Gallano A, Tordesillas-Gutierrez D, Infante J. Nonpharmacological, nonsurgical treatments for freezing of gait in Parkinson's disease: A systematic review. *Mov Disord*. 2020 Feb;35(2):204-214. doi: 10.1002/mds.27913. Epub 2019 Nov 26. PMID: 31769904.
13. Delval A, Rambour M, Tard C, et al. Freezing/festination during motor tasks in early-stage Parkinson's disease: A prospective study. *Mov Disord*. 2016;31(12):1837-1845. doi:10.1002/MDS.26762
14. Dijkstra BW, Gilat M, Cofré Lizama LE, Mancini M, Bergmans B, Verschueren SMP, Nieuwboer A. Impaired Weight-Shift Amplitude in People with Parkinson's Disease with Freezing of Gait. *J Parkinsons Dis*. 2021;11(3):1367-1380. doi: 10.3233/JPD-202370. PMID: 33749618.
15. Ehgoetz Martens KA, Shine JM, Walton CC, et al. Evidence for subtypes of freezing of gait in Parkinson's disease. *Mov Disord*. 2018;33(7):1174-1178. doi:10.1002/mds.27417
16. Espay AJ, Fasano A, Van Nuenen BFL, Payne MM, Snijders AH, Bloem BR. "On" state freezing of gait in Parkinson disease: A paradoxical levodopa-induced complication. *Neurology*. 2012;78(7):454-457. doi:10.1212/WNL.0b013e3182477ec0

17. Ferrazzoli D, Ortelli P, Madeo G, Giladi N, Petzinger GM, Frazzitta G. Basal ganglia and beyond: The interplay between motor and cognitive aspects in Parkinson's disease rehabilitation. *Neurosci Biobehav Rev.* 2018 Jul;90:294-308. doi: [10.1016/j.neubiorev.2018.05.007](https://doi.org/10.1016/j.neubiorev.2018.05.007).
18. Fietzek UM, Zwosta J, Schroeteler FE, Ziegler K, Ceballos-Baumann AO. Levodopa changes the severity of freezing in Parkinson's disease. *Parkinsonism Relat Disord.* 2013 Oct;19(10):894-6. doi: [10.1016/j.parkreldis.2013.04.004](https://doi.org/10.1016/j.parkreldis.2013.04.004). Epub 2013 May 1. PMID: 23642712.
19. Frazzitta G, Maestri R, Uccellini D, Bertotti G, Abelli P. Rehabilitation treatment of gait in patients with Parkinson's disease with freezing: A comparison between two physical therapy protocols using visual and auditory cues with or without treadmill training. *Mov Disord.* 2009;24(8):1139-1143. doi: [10.1002/mds.22491](https://doi.org/10.1002/mds.22491)
20. Frenkel-Toledo S, Giladi N, Peretz C, Herman T, Gruendlinger L, Hausdorff JM. Treadmill walking as an external pacemaker to improve gait rhythm and stability in Parkinson's disease. *Mov Disord.* 2005 Sep;20(9):1109-14. doi: [10.1002/mds.20507](https://doi.org/10.1002/mds.20507).
21. García-Ruiz PJ. Gait disturbances in Parkinson disease. Did freezing of gait exist before levodopa? Historical review. *J Neurol Sci.* 2011;307:15-17. doi: [10.1016/j.jns.2011.05.0191](https://doi.org/10.1016/j.jns.2011.05.0191).
22. Gilat M, Ginis P, Zoetewei D, et al. A systematic review on exercise and training-based interventions for freezing of gait in Parkinson's disease. *NPJ Parkinsons Dis.* 2021;7(1):81. Published 2021 Sep 10. doi: [10.1038/s41531-021-00224-4](https://doi.org/10.1038/s41531-021-00224-4)
23. Ginis P, Nackaerts E, Nieuwboer A, Heremans E. Cueing for people with Parkinson's disease with freezing of gait: A narrative review of the state-of-the-art and novel perspectives. *Ann Phys Rehabil Med* 2018;61(6):407-413. doi: [10.1016/j.rehab.2017.08.002](https://doi.org/10.1016/j.rehab.2017.08.002)
24. Ginis P, Nackaerts E, Nieuwboer A, Heremans E. Cueing for people with Parkinson's disease with freezing of gait: A narrative review of the state-of-the-art and novel perspectives. *Annals of Physical and Rehabilitation Medicine.* 2018;61(6):407-413. doi: [10.1016/j.rehab.2017.08.002](https://doi.org/10.1016/j.rehab.2017.08.002)
25. Giladi N, Nieuwboer A. Understanding and treating freezing of gait in parkinsonism, proposed working definition, and setting the stage. *Movement Disorders.* 2008;23(S2):S423-S425. doi: [10.1002/mds.21927](https://doi.org/10.1002/mds.21927)
26. Godi M, Giardini M, Nardone A, Turcato AM, Caligari M, Pisano F, Schieppati M. Curved Walking Rehabilitation with a Rotating Treadmill in Patients with Parkinson's Disease: A Proof of Concept. *Front Neurol.* 2017 Feb 28;8:53. doi: [10.3389/fneur.2017.00053](https://doi.org/10.3389/fneur.2017.00053).
27. Goetz CG, Tilley BC, Shaftman SR, et al. Movement Disorder Society-Sponsored Revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. *Mov Disord.* 2008;23(15):2129-2170. doi: [10.1002/mds.22340](https://doi.org/10.1002/mds.22340)
28. Goh, L., Canning, C. G., Song, J., Clemson, L., & Allen, N. E. (2023). The effect of rehabilitation interventions on freezing of gait in people with Parkinson's disease is unclear: a systematic review and meta-analyses. *Disability and rehabilitation, 45*(19), 3199–3218. <https://doi.org/10.1080/09638288.2022.2120099>
29. Goh L, Paul SS, Canning CG, et al. The Ziegler Test Is Reliable and Valid for Measuring Freezing of Gait in People With Parkinson Disease. *Phys Ther.* 2022;102(12):1-8. doi: [10.1093/ptj/pzac122](https://doi.org/10.1093/ptj/pzac122)
30. Heremans E, Nackaerts E, Vervoort G, et al. Amplitude manipulation evokes upper limb freezing during handwriting in patients with Parkinson's disease with freezing of gait. *PLoS One.* 2015;10(11):1-13. doi: [10.1371/journal.pone.0142874](https://doi.org/10.1371/journal.pone.0142874)
31. Herman T, Dagan M, Shema-Shiratzky S, Reches T, Brozgol M, Giladi N, Manor B, Hausdorff JM. Advantages of timing the duration of a freezing of gait-provoking test in individuals with Parkinson's disease. *J Neurol.* 2020 Sep;267(9):2582-2588. doi: [10.1007/s00415-020-09856-7](https://doi.org/10.1007/s00415-020-09856-7). Epub 2020 May 7. PMID: 32383040.
32. Herman T, Giladi N, Gruendlinger L, Hausdorff JM. Six weeks of intensive treadmill training improves gait and quality of life in patients with Parkinson's disease: a pilot study. *Arch Phys Med Rehabil.* 2007 Sep;88(9):1154-8. doi: [10.1016/j.apmr.2007.05.015](https://doi.org/10.1016/j.apmr.2007.05.015). PMID: 17826461
33. Hong M, Earhart GM. Rotating treadmill training reduces freezing in Parkinson disease: preliminary observations. *Parkinsonism Relat Disord.* 2008;14(4):359-63. doi: [10.1016/j.parkreldis.2007.07.003](https://doi.org/10.1016/j.parkreldis.2007.07.003). Epub 2007 Aug 29. PMID: 17761449; PMCID: PMC2446473.
34. Hulzinga F, Nieuwboer A, Dijkstra BW, Mancini M, Strouwen C, Bloem BR, Ginis P. The New Freezing of Gait Questionnaire: Unsuitable as an Outcome in Clinical Trials? *Mov Disord Clin Pract.* 2020 Jan 14;7(2):199-205. doi: [10.1002/mdc3.12893](https://doi.org/10.1002/mdc3.12893). PMID: 32071940; PMCID: PMC7011794.
35. Jacobs JV, Nutt JG, Carlson-Kuhta P, Stephens M, Horak FB. Knee trembling during freezing of gait represents multiple anticipatory postural adjustments. *Exp Neurol.* 2009 Feb;215(2):334-41. doi: [10.1016/j.expneurol.2008.10.019](https://doi.org/10.1016/j.expneurol.2008.10.019).

36. Jansen JAF, Capato TTC, Darweesh SKL, et al. Exploring the levodopa-paradox of freezing of gait in dopaminergic medication-naïve Parkinson's disease populations. *npj Park Dis*. 2023;9(1):130. doi:10.1038/s41531-023-00575-0
37. José Luvizutto G, Souza Silva Brito T, de Moura Neto E, Aparecida Pascucci Sande de Souza L. Altered Visual and Proprioceptive Spatial Perception in Individuals with Parkinson's Disease. *Percept Mot Skills*. 2019;127(1):98-112. doi:10.1177/0031512519880421
38. Jung SH, Hasegawa N, Mancini M, et al. Effects of the agility boot camp with cognitive challenge (ABC-C) exercise program for Parkinson's disease. *npj Park Dis*. 2020;6(1):31. doi:10.1038/s41531-020-00132-z
39. Kadivar Z, Corcos DM, Foto J, Hondzinski JM. Effect of step training and rhythmic auditory stimulation on functional performance in Parkinson patients. *Neurorehabil Neural Repair*. 2011 Sep;25(7):626-35. doi: 10.1177/1545968311401627. Epub 2011 Mar 24. PMID: 21436393.
40. Kashif M, Ahmad A, Bandpei MAM, Syed HA, Raza A, Sana V. A Randomized Controlled Trial of Motor Imagery Combined with Virtual Reality Techniques in Patients with Parkinson's Disease. *J Pers Med*. 2022;12(3):450. doi:10.3390/jpm12030450
41. King LA, Horak FB. Delaying Mobility Disability in People With Parkinson Disease Using a Sensorimotor Agility Exercise Program. *Phys Ther*. 2009;89(4):384-393. doi:10.2522/ptj.20080214
42. King LA, Mancini M, Smulders K, et al. Cognitively Challenging Agility Boot Camp Program for Freezing of Gait in Parkinson Disease. *Neurorehabilitation and Neural Repair*. 2020;34(5):417-427. doi:10.1177/1545968320909331
43. King LA, Peterson DS, Mancini M, et al. Do cognitive measures and brain circuitry predict outcomes of exercise in Parkinson Disease: A randomized clinical trial. *BMC Neurol*. 2015;15(1):4-11. doi:10.1186/s12883-015-0474-2
44. King LA, Priest KC, Nutt J, et al. Comorbidity and Functional Mobility in Persons with Parkinson Disease. *Arch Phys Med Rehabil*. 2014;95(11):2152-2157. doi:10.1016/j.apmr.2014.07.396
45. Koehler PJ, Nonnekes J, Bloem BR. Freezing of gait before the introduction of levodopa. *Lancet Neurol*. 2021 Feb;20(2):97. doi: 10.1016/S1474-4422(19)30091-2. Epub 2019 Apr 11. PMID: 30982582.
46. Kwok JYY, Smith R, Chan LML, Lam LCC, Fong DYT, Choi EPH, Lok KYW, Lee JJ, Auyeung M, Bloem BR. Managing freezing of gait in Parkinson's disease: a systematic review and network meta-analysis. *J Neurol*. 2022 Jun;269(6):3310-3324. doi: 10.1007/s00415-022-11031-z.
47. Leung IHK, Walton CC, Hallock H, Lewis SJG, Valenzuela M, Lampit A. Cognitive training in Parkinson disease: A systematic review and meta-analysis. *Neurology*. 2015;85(21):1843-1851. doi:10.1212/WNL.0000000000002145
48. Lewis S, Factor S, Giladi N, Nieuwboer A, Nutt J, Hallett M. Stepping up to meet the challenge of freezing of gait in Parkinson's disease. *Transl Neurodegener*. 2022;11(1):23. doi:10.1186/s40035-022-00298-x
49. Maidan I, Nieuwhof F, Bernad-Elazari H, Bloem BR, Giladi N, Hausdorff JM, et al. Evidence for Differential Effects of 2 Forms of Exercise on Prefrontal Plasticity During Walking in Parkinson's Disease. *Neurorehabil Neural Repair*. 2018 Mar;32(3):200-208. doi: 10.1177/1545968318763750.
50. Maidan I, Rosenberg-Katz K, Jacob Y, Giladi N, Hausdorff JM, Mirelman A. Disparate effects of training on brain activation in Parkinson disease. *Neurology*. 2017 Oct 24;89(17):1804-1810. doi: 10.1212/WNL.0000000000004576.
51. Martens KAE, Silveira CRA, Intzandt BN, Almeida QJ. Overload From Anxiety: A Non-Motor Cause for Gait Impairments in Parkinson's Disease. *J Neuropsychiatry Clin Neurosci*. 2018;30:77-80. doi:10.1176/appi.neuropsych.16110298
52. Mori L, Putzolu M, Bonassi G, Galeoto G, Mezzarobba S, Trompetto C, Avanzino L, Marchese R, Abbruzzese G, Pelosin E. Haptic perception of verticality correlates with postural and balance deficits in patients with Parkinson's disease. *Parkinsonism Relat Disord*. 2019 Sep;66:45-50. doi: 10.1016/j.parkreldis.2019.06.026. Epub 2019 Jul 2. PMID: 31279636.
53. Nieuwboer A, Giladi N. Characterizing freezing of gait in Parkinson's disease: Models of an episodic phenomenon. *Mov Disord*. 2013;28(11):1509-1519. doi:10.1002/mds.25683
54. Nieuwboer A, Rochester L, Herman T, et al. Reliability of the new freezing of gait questionnaire: Agreement between patients with Parkinson's disease and their carers. *Gait Posture*. 2009;30(4):459-463. doi:10.1016/j.gaitpost.2009.07.108
55. Nóbrega LR, Rocon E, Pereira AA, Andrade A de O. A Novel Physical Mobility Task to Assess Freezers in Parkinson's Disease. *Healthcare*. 2023;11(3):409. doi:10.3390/healthcare11030409
56. Nonnekes J, Giladi N, Guha A, Fietzek UM, Bloem BR, Růžička E. Gait festination in parkinsonism: introduction of two phenotypes. *J Neurol*. 2019;266(2):426-430. doi:10.1007/s00415-018-9146-7

57. Nonnekes J, Snijders AH, Nutt JG, Deuschl G, Giladi N, Bloem BR. Freezing of gait: a practical approach to management. *Lancet Neurol*. 2015;14(7):768-778. doi:10.1016/S1474-4422(15)00041-1
58. Nonnekes J, Nieuwboer A. Towards Personalized Rehabilitation for Gait Impairments in Parkinson's Disease. Brundin P, Langston JW, Bloem BR, eds. *Journal of Parkinson's Disease*. 2018;8(s1):S101-S106. doi:[10.3233/JPD-181464](https://doi.org/10.3233/JPD-181464)
59. Nutt JG, Bloem BR, Giladi N, Hallett M, Horak FB, Nieuwboer A. Freezing of gait: moving forward on a mysterious clinical phenomenon. *Lancet Neurol*. 2011;10(8):734-744. doi:10.1016/S1474-4422(11)70143-0
60. Pelosin E, Avanzino L, Bove M, Stramesi P, Nieuwboer A, Abbruzzese G. Action observation improves freezing of gait in patients with Parkinson's disease. *Neurorehabil Neural Repair*. 2010 Oct;24(8):746-52. doi: 10.1177/1545968310368685. Epub 2010 May 7. PMID: 20453155.
61. Pelosin E, Barella R, Bet C, Magioncalda E, Putzolu M, Di Biasio F, Cerulli C, Casaleggio M, Abbruzzese G, Avanzino L. Effect of Group-Based Rehabilitation Combining Action Observation with Physiotherapy on Freezing of Gait in Parkinson's Disease. *Neural Plast*. 2018 May 27;2018:4897276. doi: 10.1155/2018/4897276. PMID: 29977280; PMCID: PMC5994277.
62. Peterson DS, King LA, Cohen RG, Horak FB. Cognitive Contributions to Freezing of Gait in Parkinson Disease: Implications for Physical Rehabilitation. *Physical Therapy*. 2016;96(5):659-670. doi:[10.2522/ptj.20140603](https://doi.org/10.2522/ptj.20140603)
63. Rafferty MR, Nettnin E, Goldman JG, MacDonald J. Frameworks for Parkinson's Disease Rehabilitation Addressing When, What, and How. *Curr Neurol Neurosci Rep*. 2021;21(3):12. doi:10.1007/s11910-021-01096-0
64. Rutz DG, Benninger DH. Physical Therapy for Freezing of Gait and Gait Impairments in Parkinson Disease: A Systematic Review. *PM&R*. 2020 Nov;12(11):1140-1156. doi: 10.1002/pmrj.12337. Epub 2020 May 19.
65. Sarasso E, Agosta F, Piramide N, Gardoni A, Canu E, Leocadi M, Castelnovo V, Basaia S, Tettamanti A, Volontè MA, Filippi M. Action Observation and Motor Imagery Improve Dual Task in Parkinson's Disease: A Clinical/fMRI Study. *Mov Disord*. 2021 Nov;36(11):2569-2582. doi: 10.1002/mds.28717. Epub 2021 Jul 19. PMID: 34286884
66. Shah V V., Vitorio R, Hasegawa N, et al. Effects of a Cognitively Challenging Agility Boot Camp Program (ABC-C) on Balance and Gait in People With Parkinson's Disease: Does Freezing of Gait Status Matter? *Neurorehabil Neural Repair*. 2022;36(9):603-612. doi:10.1177/15459683221119757
67. Silva-Batista C, Corcos DM, Kanegusuku H, et al. Balance and fear of falling in subjects with Parkinson's disease is improved after exercises with motor complexity. *Gait Posture*. 2018;61(May 2017):90-97. doi:10.1016/j.gaitpost.2017.12.027
68. Silva-Batista C, Corcos DM, Roschel H, Kanegusuku H, Gobbi LT, Piemonte ME, Mattos EC, DE Mello MT, Forjaz CL, Tricoli V, Ugrinowitsch C. Resistance Training with Instability for Patients with Parkinson's Disease. *Med Sci Sports Exerc*. 2016 Sep;48(9):1678-87. doi: 10.1249/MSS.0000000000000945.
69. Silva-Batista C, Lima-Pardini AC, Nucci MP, et al. A Randomized, Controlled Trial of Exercise for Parkinsonian Individuals With Freezing of Gait. *Mov Disord*. 2020;35(9):1607-1617. doi:10.1002/mds.28128
70. Snijders AH, Haaxma CA, Hagen YJ, Munneke M, Bloem BR. Freezer or non-freezer: Clinical assessment of freezing of gait. *Park Relat Disord*. 2012;18(2):149-154. doi:10.1016/j.parkreldis.2011.09.006
71. Snijders AH, Nijkrake MJ, Bakker M, Munneke M, Wind C, Bloem BR. Clinimetrics of freezing of gait. *Mov Disord*. 2008;23(S2):S468-S474. doi:10.1002/mds.22144
72. Sozzi S, Schieppati M. Stepping in Place While Voluntarily Turning Around Produces a Long-Lasting Posteffect Consisting in Inadvertent Turning While Stepping Eyes Closed. *Neural Plast*. 2016;2016:7123609. doi: 10.1155/2016/7123609.
73. Taghizadeh G, Fereshted SM, Martinez-Martin P, Joghataei MT, Mahdizadeh F, Sabbaghi S, Goudarzi S, Meimandi M, Habibi SAH, Mehdizadeh M. Clinimetrics of the Freezing of Gait Questionnaire for Parkinson Disease During the "off" State. *Basic Clin Neurosci*. 2021 Jan-Feb;12(1):69-78. doi: 10.32598/bcn.12.1.882.11. Epub 2021 Jan 1. PMID: 33995929; PMCID: PMC8114865.
74. Tosserams A, Bloem BR, Ehgoetz Martens KA, Helmich RC, Kessels RPC, Shine JM, Taylor NL, Wainstein G, Lewis SJG, Nonnekes J. Modulating arousal to overcome gait impairments in Parkinson's disease: how the noradrenergic system may act as a double-edged sword. *Transl Neurodegener*. 2023 Mar 26;12(1):15. doi: 10.1186/s40035-023-00347-z. PMID: 36967402; PMCID: PMC10040128.

75. Tosserams A, Nijkrake MJ, Sturkenboom IHWM, Bloem BR, Nonnekes J. Perceptions of Compensation Strategies for Gait Impairments in Parkinson's Disease: A Survey Among 320 Healthcare Professionals. *J Parkinsons Dis.* 2020;10:1775-1778. doi:10.3233/JPD-202176
76. Tosserams A, Nonnekes J. How I Examine My Patient: A Practical Guide to the Evaluation of Compensation Strategies for Gait Impairment in Parkinson's Disease. *J Parkinsons Dis.* 2022;12:2005-2008. doi:10.3233/JPD-223296
77. Tseng, Ing-Jy MSN; Yuan, Rey-Yue MD; Jeng, Chii PhD. Treadmill Training Improves Forward and Backward Gait in Early Parkinson Disease. *American Journal of Physical Medicine & Rehabilitation* 94(10):p 811-819, October 2015. | DOI: 10.1097/PHM.0000000000000273
78. Vieira-Yano B, Martini DN, Horak FB, et al. The Adapted Resistance Training with Instability Randomized Controlled Trial for Gait Automaticity. *Mov Disord.* 2021;36(1):152-163. doi:10.1002/mds.28298
79. Walton CC, Mowszowski L, Gilat M, Hall JM, O'Callaghan C, Muller AJ, Georgiades M, Szeto JYY, Ehgoetz Martens KA, Shine JM, Naismith SL, Lewis SJG. Cognitive training for freezing of gait in Parkinson's disease: a randomized controlled trial. *NPJ Parkinsons Dis.* 2018 May 18;4:15. doi: 10.1038/s41531-018-0052-6.
80. Weiss D, Schoellmann A, Fox MD, et al. Freezing of gait: understanding the complexity of an enigmatic phenomenon. *Brain.* 2020;143(1):14-30. doi:10.1093/brain/awz314
81. Zhang LL, Canning SD, Wang XP. Freezing of Gait in Parkinsonism and its Potential Drug Treatment. *Curr Neuropharmacol.* 2016;14(4):302-6. doi: 10.2174/1570159x14666151201190040. PMID: 26635194; PMCID: PMC4876585.
82. Ziegler, K., Schroeteler, F., Ceballos-Baumann, A.O. and Fietzek, U.M. (2010), A new rating instrument to assess festination and freezing gait in Parkinsonian patients. *Mov. Disord.*, 25: 1012-1018. <https://doi.org/10.1002/mds.22993>

PWR! uses the latest research to inform our programs, workshops and resources. The full body of research referenced during the workshop is updated regularly and can be viewed at:

pwr4life.org/Parkinson-research